Takeshi KAWAZOE et al. Serial No. 10/528,150 Attorney Docket No. 2005_0459A September 15, 2008

AMENDMENTS TO THE CLAIMS

1-2. (Cancelled)

3. (Currently amended) A method for inhibition of keloid and/or hypertrophic scar formation in a course of therapy of wound or dermal injury and for inhibition treatment of pain or pruritus on the a wound lesion of keloid and/or hypertrophic scar—which comprises administering a medicament containing acetylsalicylic acid or its pharmacologically acceptable salt-in an effective amount to said wound lesion of a patient.

4. (Cancelled)

- 5. (Currently amended) A method for inhibition of keloid and/or hypertrophic scar formation in a course of therapy of wound or dermal injury which comprises administering a medicament containing acetylsalicylic acid or its pharmacologically acceptable salt-in an effective amount to a wound lesion of a patient.
- 6. (New) A method for inhibition of keloid and/or hypertrophic scar formation in a course of therapy of wound or dermal injury which comprises administering a medicament containing acetylsalicylic acid in an effective amount of 0.005 to 1000 μ g/g per tissue weight to a wound lesion of a patient.
- 7. (New) A method for inhibition of keloid and/or hypertrophic scar formation in a course of therapy of wound or dermal injury which comprises administering a medicament containing acetylsalicylic acid in an effect amount of 0.01 to 800 μ g/g per tissue weight to a wound lesion of a patient.